

Basic Guidelines to Animal Homeopathic Dosing

in accordance with Classical Homeopathic Prescribing Principles

Handling and Storage of Homeopathic Remedies:

Dispense remedies into the cap of the vial and not onto your hand, as in doing so, you may weaken the energy field in them. Never leave remedies in direct sunlight. Never store remedies on or near an electromagnetic force (such as TV, radio, microwave, electric blanket, etc.). Never open or store remedies in the presence of a strong odor, i.e.. camphorated products, nail polish remover, toothpaste, oil based paints, etc.

Acute Care: Dose every 20 minutes to 1 hour for 3-4 hours. Decrease frequency as symptoms improve.

If no improvement is seen, you may have the wrong remedy - check your symptom picture again to choose another remedy. If improvement is seen but fails to hold, go to 200C.

Dosing: Medium and Low Potencies (6C, 12C, 30C, 200C) 2-3 pills per dose is all that's needed. The more severe the symptoms, the more frequent the dose is given in Acute Care First Aid Ailments.

The higher the potency, the less frequently you give the remedy. Always adjust remedy frequency up or down according to the symptoms.

Never leave a remedy that's working!

Base Frequency of Doses According to Symptom Severity!

Dispensing Remedy: Dispense remedy into cap of vial, and then to one of the methods below.

Direct Dosing: The easiest way to administer remedies to an animal if they will take things from your hand. Animals who have never had remedies before may refuse them at first, but once they realize what they are, will willingly take them in the future.

In Food: Give one dose in meals or treats at recommended intervals.

Note: For animals who will not take a remedy from hand, will not drink or are stressed by the water potencies in syringe. The least preferred method, but it will work.

Water Potency: Best used when an animal needs a remedy and you can't be there to administer it at recommended intervals, or if they are unconscious, or unwilling to take pellets from your hand. For use with lower to medium potencies only, and for limited time.

1. Put one dose (2-3 pellets) into their water bowl and fill with clean water.

Note: Give a second water bowl so that they may choose to take the remedy or not. They will dose themselves accordingly, in most cases. It is best to be sure they have the first dose or drink before leaving them to their own dosing.

2. Put one dose in 1/2 glass clean water and stir briskly with a stainless steel spoon. Use a veterinary syringe (without needle) to suck up a small amount of water remedy, then administer to animal. Refrigerate unused remedy with a lid, stirring briskly before next dosage is extracted with syringe. Label syringe with Remedy name and potency. Do not use for any other remedy.

Note: Unconscious animals can NOT swallow!

Avoid choking by only placing 1 or 2 drops of the remedy in their mouth between lip and gums.

For Severe or Life Threatening Injuries

SEEK PROFESSIONAL, EMERGENCY CARE IMMEDIATELY!

Use these first aid remedies while transporting to, or while waiting for emergency medical personnel to arrive at the scene.



© 1999 Animal Wellness. All Rights Reserved Worldwide.